

UC Davis

Dermatology Online Journal

Title

Fish odor syndrome: a case report of trimethylaminuria

Permalink

<https://escholarship.org/uc/item/9550x15j>

Journal

Dermatology Online Journal, 20(1)

Authors

Ulman, Catherine A
Trevino, Julian J
Miller, Marvin
et al.

Publication Date

2014

DOI

10.5070/D3201021260

Copyright Information

Copyright 2014 by the author(s). This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Letter

Fish odor syndrome: a case report of trimethylaminuria

Catherine A. Ulman BA¹, Julian J. Trevino MD², Marvin Miller MD³, Rishi K. Gandhi MD²

Dermatology Online Journal 20 (1): 19

Wright State University, ¹Boonshoft School of Medicine, ²Department of Dermatology, Dayton, OH

³Dayton Children's Medical Center, Department of Medical Genetics, Dayton, OH

Correspondence:

Rishi K. Gandhi, M.D.
725 University Blvd.
Dayton, OH 45435
rishi.gandhi@wright.edu

Abstract

Trimethylaminuria is a rare, autosomal recessive, metabolic disorder that results in accumulation of trimethylamine (TMA), which smells like rotten fish. The chemical is excreted in sweat and urine owing to a deficiency in the enzyme flavin monooxygenase 3 (FMO3). We report a case of trimethylaminuria in a 12-year-old girl. The patient failed treatment with diet and hygiene modification, but achieved symptomatic improvement after a four-month course of metronidazole.

Keywords: Trimethylaminuria, Fish Odor Syndrome, Metabolic Disorders

Case synopsis

A 12-year-old girl was evaluated in the dermatology clinic for a chronic foul-smelling body odor. The patient's past medical history was significant for anxiety with generalized hyperhidrosis. Her symptoms flared during periods of anxiety and failed to improve with use of scented antiperspirant deodorants and frequent bathing. The patient was ostracized and severely ridiculed at school.

The patient was adopted, and her biologic family history was unknown. Physical examination was unremarkable except for noticeable foul-smelling odor.

The diagnosis of trimethylaminuria was confirmed by calculating the percent trimethylamine-N-oxide (TMAO) in the patient's urine. The patient's trimethylamine level was 80.7 $\mu\text{mole}/\text{mmole}$ creatinine and the level of trimethylamine-N-oxide was 0.6 $\mu\text{mole}/\text{mmole}$ creatinine from which the percent trimethylamine-N-oxide was calculated to be 0.7% (normal is considered >92%).

The patient experienced mild improvement on a choline, tyramine, and indole-restricted diet. Significant reduction in the odor was achieved after a four-month course of metronidazole 500 mg twice daily.

Discussion

Trimethylaminuria is a rare metabolic disorder resulting in accumulation of trimethylamine (TMA) in sweat and urine owing to enzymatic deficiency of flavin monooxygenase 3 (FMO3) caused by mutations in the FMO3 gene. Primary trimethylaminuria has an autosomal recessive inheritance pattern [1].

Normally, hepatic FMO3 oxidizes TMA, which smells like rotting fish, into trimethylamine-N-oxide (TMAO), which is odorless [1]. In trimethylaminuria, TMA is excreted through the sweat and urine, resulting in a characteristic fishy body odor [2]. Increased sweating related to exercise or anxiety exacerbates the odor [3].

Initial treatment includes restriction from foods containing choline, lecithin, and carnitine (saltwater fish, liver, egg yolk, soy beans, and legumes), which are precursors to TMA [1, 2]. The excessive consumption of milk has also been linked to increased TMA excretion [4]. Even with rigorous adherence to the restricted diet, the amount of TMA excreted fluctuates [3]. Frequent bathing and deodorants may help conceal the odor [5].

If clinical improvement is not accomplished through diet alone, metronidazole may be used to decrease microbial gut flora and reduce TMA production [2, 3, 5]. Neomycin, activated charcoal, and lactulose have been used in the past, but frequently cause diarrhea. To prevent treatment failure related to drug resistance, the administration of metronidazole for one- to two-week intervals interspersed with drug-free periods is advised [3].

References

1. Chalmers RA, Bain MD, Michelakakis H, Zschocke J, Iles RA. Diagnosis and management of trimethylaminuria (FMO3 deficiency) in children. *J Inherit Metab Dis*. 2006 Feb;29:162-172. [PMID: 16601883]
2. Mackay RJ, McEntyre CJ, Henderson C, Lever M, George PM. Trimethylaminuria: causes and diagnosis of a socially distressing condition. *Clin Biochem Rev*. 2011 Feb;31:33-39. [PMID: 21451776]
3. Treacy E, Johnson D, Pitt JJ, Danks DM. Trimethylaminuria, fish odour syndrome: a new method of detection and response to treatment with metronidazole. *J Inherit Metab Dis*. 1995 Feb;18:306-312. [PMID: 7474897]
4. Rothschild JG, Hansen RC. Fish odor syndrome: trimethylaminuria with milk as chief dietary factor. *Pediatr Dermatol*. 1985 Nov;3:38-39. [PMID: 4070086]
5. Ferrari ND, Nield LS. Smelling like dead fish: a case of trimethylaminuria in an adolescent. *Clin Pediatrics*. 2006 Nov;45:864-866. [PMID: 17041178]