

UC Davis

Dermatology Online Journal

Title

Risk factors for facial telangiectasia

Permalink

<https://escholarship.org/uc/item/9kh9g4t5>

Journal

Dermatology Online Journal, 22(9)

Authors

Hamer, Merel A
Pardo, Luba M
Gunn, David A
[et al.](#)

Publication Date

2016

DOI

10.5070/D3229032561

Copyright Information

Copyright 2016 by the author(s). This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Peer reviewed

Abstract

Risk factors for facial telangiectasia

Merel A. Hamer¹, Luba M. Pardo¹, David A. Gunn², Tamar Nijsten¹

Dermatology Online Journal 22 (9)

¹Department of Dermatology, Erasmus MC University Medical Center Rotterdam, The Netherlands

²Unilever Research and Development, Colworth Science Park, Sharnbrook, Bedfordshire, United Kingdom

Background

Telangiectasia or red veins are one of the main features of facial skin aging. To date there are few studies investigating risk factors for telangiectasia. We investigated environmental risk factors in a population-based cohort study, using a digital continuous outcome measure.

Methods

Telangiectasia were quantified digitally from standardized 3-dimensional facial photographs of 2886 North-European participants (56.8% female, median age 66.9) from the Rotterdam Study, a prospective population-based cohort study. Age- and sex adjusted as well as fully adjusted multivariable linear regressions were performed to investigate associations between potential environmental factors and the amount of telangiectasia. Men and women were analyzed separately.

Results

The mean facial area covered by telangiectasia was higher in women (median area 0.95%, interquartile range 0.62- 1.4) than in men (median area 0.76%, interquartile range 0.48-1.2). Besides age (men $\beta=0.016$, P-value<0.001; women $\beta=0.016$, P-value<0.001), smoking (men $\beta=0.29$, P-value<0.001; women $\beta=0.32$, P-value<0.001), a high susceptibility to sunburn (men $\beta=0.082$, P-value<0.001; women $\beta=0.076$, P-value<0.001) and light skin color (pale against olive-colored skin in men $\beta=0.26$, P-value<0.001; in women $\beta=0.23$, P-value<0.001) were independent significant factors contributing to telangiectasia for both sexes. Additionally in men, a lower education also showed an association (medium vs. high $\beta=0.062$, P-value=0.008). Alcohol showed a negative association with telangiectasia ($\beta=-0.028$, P-value<0.001). In women, a higher free androgen index was associated with less telangiectasia ($\beta=-0.0053$, P-value<0.001).

Conclusion

In this large cohort study of facial telangiectatic photoaging, environmental factors associated with telangiectasia are described, implicating possible new prevention strategies for this form of skin aging.